

Table S1. Details of fungal coinfections in included studies

Study	<i>Aspergillus</i> (%)	<i>Candida</i> (%)	Other fungi (%)
Zhu et al. [21]	<i>Aspergillus</i> : 23.35	<i>Candida</i> : 0.78	<i>Cryptococcus</i> : 0.39
Agrifoglio et al. [22]	<i>Aspergillus fumigatus</i> complex: 1.44	<i>Candida</i> sp.: 4.32 <i>C. albicans</i> : 6.47 <i>C. parapsilosis</i> : 2.88 <i>C. glabrata</i> : 1.44	-
Chowdhary et al. [23]	-	<i>C. auris</i> : 1.68 <i>C. albicans</i> : 0.5 <i>C. tropicalis</i> : 0.17 <i>C. krusei</i> : 0.17	-
Intra et al. [25]	<i>Aspergillus fumigatus</i> : 1.64	<i>Candida albicans</i> : 22.95 <i>Candida glabrata</i> : 6.56	-
Segrelles-Calvo et al. [26]	<i>Aspergillus</i> spp: 3.26 <i>A. nige</i> : 0.93 <i>A. flavus</i> : 0.93 <i>A. fumigatus</i> : 1.4	-	-
Wang et al. [27]	<i>Aspergillus</i> : 7.69	-	-

-, not applicable.