

Table S2. Adverse pregnancy outcomes according to gestational weight gain from second to third trimester in pregnant women who were prepregnancy nonobese

Variable	Total (n = 166)	Insufficient GWG (n = 92)	Appropriate GWG (n = 48)	Excessive GWG (n = 26)	p-value
Any adverse pregnancy outcomes ^{a)}	90 (54.2)	47 (51.1)	25 (52.1)	18 (69.2)	0.245
Maternal					
Any adverse maternal outcomes ^{a)}	73 (44.0)	36 (39.1)	21 (43.8)	16 (61.5)	0.127
HDP	5 (3.0)	0 (0)	3 (6.3)	2 (7.7)	0.018
Preterm birth	13 (7.8)	7 (7.6)	4 (8.3)	2 (7.7)	1.000
Cesarean delivery	59 (35.5)	27 (29.3)	17 (35.4)	15 (57.7)	0.029
Delivery complications	15 (9.0)	11 (12.0)	1 (2.1)	3 (11.5)	0.100
Infant					
Any adverse infant outcomes ^{a)}	37 (22.3)	19 (20.7)	12 (25.0)	6 (23.1)	0.837
Low birth weight	5 (3.0)	4 (4.3)	1 (2.1)	0 (0.0)	0.698
High birth weight	16 (9.6)	6 (6.5)	5 (10.4)	5 (19.2)	0.142
NICU admission	15 (9.0)	9 (9.8)	5 (10.4)	1 (3.8)	0.659
Congenital anomaly	5 (3.0)	3 (3.3)	2 (4.2)	0 (0)	0.841

Data are presented as n (%).

GWG, gestational weight gain; HDP, hypertensive disorders of pregnancy; NICU, neonatal intensive care unit.

^{a)}Any of the variables below.