

Table S3. Adverse pregnancy outcomes according to gestational weight gain from second to third trimester in pregnant women who were prepregnancy obese

Variable	Total (n = 69)	Insufficient GWG (n = 31)	Appropriate GWG (n = 20)	Excessive GWG (n = 18)	p-value
Any adverse pregnancy outcomes ^{a)}	50 (72.5)	21 (67.7)	16 (80.0)	13 (72.2)	0.633
Maternal					
Any adverse maternal outcomes ^{a)}	47 (68.1)	19 (61.3)	15 (75.0)	13 (72.2)	0.538
HDP ^{b)}	2 (3.3)	2 (0.0)	0 (0)	0 (0)	0.493
Preterm birth	4 (5.8)	2 (6.5)	0 (0)	2 (11.1)	0.274
Cesarean delivery	42 (60.9)	18 (58.1)	14 (70.0)	10 (55.6)	0.602
Delivery complications	9 (13.0)	2 (6.5)	4 (20.0)	3 (16.7)	0.274
Infant					
Any adverse infant outcomes ^{a)}	13 (18.8)	5 (16.1)	5 (25.0)	3 (16.7)	0.730
Low birth weight	0 (0)	0 (0)	0 (0)	0 (0)	-
High birth weight	6 (8.7)	1 (3.2)	3 (15.0)	2 (11.1)	0.357
NICU admission	8 (11.6)	4 (12.9)	3 (15.0)	1 (5.6)	0.714
Congenital anomaly	1 (1.4)	0 (0)	1 (5.0)	0 (0)	0.551

Data are presented as n (%).

GWG, gestational weight gain; HDP, hypertensive disorders of pregnancy; NICU, neonatal intensive care unit.

^{a)}Any of the variables below. ^{b)}Only 61 participants were included, excluding those with a history of hypertension or no information on the history of hypertension.